

This schedule is recommended as a general guideline for product usage. Take local biotic and abiotic conditions into account when using this schedule.



ROOTING PHASE

GROWTH PHASE

EARLY BLOOM PHASE

BLOOM PHASE

RIPENING PHASE



COCO MAX

A: 1 - 2 ml/L
B: 1 - 2 ml/L

A: 3 - 4 ml/L
B: 3 - 4 ml/L

A: 3 - 4 ml/L
B: 3 - 4 ml/L

A: 3 - 4 ml/L
B: 3 - 4 ml/L

A: 3 - 4 ml/L
B: 3 - 4 ml/L

ROOTBASTIC



0,1 - 0,3 ml/L



ATAZYME

1 - 3 ml/L

1 - 3 ml/L

1 - 3 ml/L

1 - 3 ml/L

BLOOMBASTIC



0,5 - 1 ml/L

0,5 - 1 ml/L

0,5 - 1 ml/L



CALMAG

0,1 - 1 ml/L

0,1 - 1 ml/L

0,1 - 1 ml/L

0,1 - 1 ml/L

0,1 - 1 ml/L

B'CUZZ SILIC BOOST



0,1 ml/L

0,1 ml/L

0,1 ml/L

0,1 ml/L

0,1 ml/L